Enbumyst™
(bumetanide nasal spray)
0.5 mg per device
INSTRUCTIONS FOR USE

This Instructions for Use contains Information on how to use ENBUMYST

- Talk to your healthcare provider on how to use **ENBUMYST**.
- Read this Instructions for Use before using **ENBUMYST**.

Storing ENBUMYST

- Store in carton at room temperature of 59°F to 77°F (15°C to 25°C).
- Do not freeze.
- Keep **ENBUMYST** and all medicines out of the reach of children.


Important Information You Need to Know Before Using ENBUMYST

- For use in the nose only.
- Each blister contains 1 device. Do not remove from blister until ready to use.
- Check the expiration date. Do not use past the expiration date on the blister or if the device appears damaged.
- Do not test, prime (pre-spray), or reuse the device. Each device can only be sprayed 1 time.
- Refer to the table and only remove the number of devices needed for your prescribed dose (up to 4).

Dose	Number of Nasal Spray Devices
0.5 mg	1 device
1 mg	2 devices
1.5 mg	3 devices
2 mg	4 devices

Step 1

Remove Device from Blister



Before first device use only, blow your nose gently to clear both nostrils.

Remove the number of blisters needed for your prescribed dose from the carton.

Peel back the tab with the triangle to open the blister and remove the device.

Step 2

Properly Hold Device



Hold **ENBUMYST** with your thumb on the bottom of the plunger and your index and middle fingers on either side of the nozzle.

Do not press the plunger yet.

Step 3

Insert Device into Nose




Tilt your head back slightly.

Gently insert the tip of the nozzle into one nostril until your fingers on either side of the nozzle are against the bottom of your nose.

Step 4

Press Plunger Firmly



Gently press the other nostril with your finger to keep it closed.

Make sure that the tip of the device is **pointed straight into the nose** and **not** against the side (wall) of the nose.

Slowly breath through the open nostril.

Press the plunger firmly to give the dose of **ENBUMYST**.

Step 5

Remove and Dispose of Device



Remove **ENBUMYST** nasal spray from the nostril after giving the dose.

Dispose of (throw away) the used device and blister in the household trash.

Dose Completion



Alternate Nostrils

If your prescribed dose requires more than 1 ENBUMYST device, **repeat Steps 1 to 5** to complete your dose. **Alternate each time** between your right and left nostrils.

Note: Do not blow your nose. Keep your head tilted back for approximately 1 minute before using another device. This will allow the medicine to go into your nasal passage without dripping out.

For more information visit ENBUMYST.com or call 1-877-300-5339

This Instructions for Use has been approved by the U.S. Food and Drug Administration. 09/2025

©2025 Corstasis Therapeutics Inc. All Rights Reserved. **ENBUMYST** is a registered trademark of Corstasis Therapeutics Inc.