Enbumyst®
(bumetanide nasal spray) 0.5 mg

The only FDA-approved nasal spray diuretic

It's Not Oral.
It's Not IV.
It's ENBUMYST.

A short-term diuretic option for at-home use.



Not actual size.

What is ENBUMYST?

ENBUMYST is a prescription medicine used in adults to treat swelling caused by excess fluid in the body (edema) from congestive heart failure, liver disease, and kidney disease.

What is the most important information I should know about ENBUMYST?

- **ENBUMYST is a diuretic** that helps your body remove fluid. Losing too much fluid can cause dehydration, low blood volume, and increase the risk of blood clots. Blood abnormalities may include decreased salts (electrolytes), and increased nitrogen, glucose and uric acid. The risk of these abnormalities is higher in elderly people, those using higher doses, or those who do not get enough electrolytes by mouth.

Please see Important Safety Information throughout and full Prescribing Information and Patient Information for ENBUMYST.

Enbumyst™
(bumetanide nasal spray) 0.5 mg

Why might my healthcare provider prescribe ENBUMYST?

You may be prescribed ENBUMYST to treat swelling (edema) related to heart failure, liver disease, or kidney disease.

This may include situations where:

- Swelling or fluid buildup (edema) is worsening.
- Oral diuretic pills are not the right choice.
- Absorption of oral medications may be affected.
- Short-term treatment of excess fluid is needed during recovery as you transition home.

ENBUMYST is not intended for long-term (chronic) use. It is designed to manage swelling due to too much fluid (edema), with the goal of transitioning to oral diuretic therapy when appropriate.

Actor portrayal

IMPORTANT SAFETY INFORMATION (cont'd)

- **You should not use ENBUMYST if you have severe electrolyte loss.**
  Tell your healthcare provider if you experience unusual thirst; dizziness or lightheadedness; weakness or tiredness; fast heart rate; dry mouth or tongue; sudden pain; redness or swelling in your joints; pain, swelling, redness, or warmth in your leg; shortness of breath or fast breathing; high blood sugar; dark urine; pale, tight, cool or clammy skin; muscle cramps; confusion.

Do not use ENBUMYST if you have low or no urine production or are allergic to bumetanide or to any of the ingredients in ENBUMYST.

Please see Important Safety Information throughout and full Prescribing Information and Patient Information for ENBUMYST.

Key Steps for Administration



Step 1: PREPARE – Remove device from blister; blow nose before first device use only.

Step 2: HOLD device with thumb on plunger.



Step 3: INSERT device straight into nostril while tilting head back.



Step 4: CLOSE the other nostril, breathe in slowly, and press firmly.

Step 5: REMOVE device from nostril and discard.

If your prescribed dose requires more than 1 device: Repeat steps 1–5 while alternating nostrils. Keep head tilted back for about 1 minute before using another device.

Read the complete Instructions for Use before administering ENBUMYST (bumetanide nasal spray.) Scan the QR code to view full instructions including a demonstration video.

IMPORTANT SAFETY INFORMATION (cont'd)

Before using ENBUMYST, tell your healthcare provider if you have low electrolytes, are on a low-salt diet, have diarrhea, diabetes, nasal problems such as a runny or stuffy nose or polyps, a history of nasal injury or surgery, are or plan to become pregnant, are or plan to breastfeed.

Tell your healthcare provider about all the medicines you take, including prescriptions, over-the-counter medicines, vitamins, and herbal supplements.

ENBUMYST may cause serious side effects, such as worsening kidney function and ringing in the ears or hearing loss.

Please see Important Safety Information throughout and full Prescribing Information and Patient Information for ENBUMYST.

Enbumyst
(bumetanide nasal spray) 0.5 mg

Common Questions

How long will I use ENBUMYST?

ENBUMYST is for short-term use, based on your healthcare provider's instructions. It is not intended for long-term (chronic) use.

How many devices should I use?

Use exactly the number of devices prescribed by your healthcare provider. Each device delivers one spray and is for one-time use only.

How should I dispose of used or expired devices?

After use, discard the device and blister in the household trash. For expired product, ask your pharmacist about drug take-back options.

How should I store ENBUMYST?

Store at room temperature (59°F–77°F / 15°C–25°C). Do not freeze.

Learn more at ENBUMYST.com

IMPORTANT SAFETY INFORMATION (cont'd)

The most common side effects of ENBUMYST include dehydration, headache, muscle cramps, dizziness, low blood pressure, nausea, and abnormal brain functioning with certain liver problems.

These are not all the possible side effects of ENBUMYST. Talk to your healthcare provider for more information.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see Important Safety Information throughout and full Prescribing Information and Patient Information for ENBUMYST.

All trademarks and trade names are the property of their respective owners. © 2026 Esperion Therapeutics, Inc. All rights reserved. US-ENBU-2600074-2

ESPERION